

Asymptotic analysis of the Askey-scheme II: from Charlier to Hermite

Diego Dominici *

Department of Mathematics
State University of New York at New Paltz
75 S. Manheim Blvd. Suite 9
New Paltz, NY 12561-2443
USA

Phone: (845) 257-2607

Fax: (845) 257-3571

February 2, 2008

Abstract

We analyze the Hermite polynomials $H_n(\xi)$ and their zeros asymptotically as $n \rightarrow \infty$, using the limit relation between the Charlier and Hermite polynomials. Our formulas involve some special functions and they yield very accurate approximations.

Keywords: Hermite polynomials, Askey-scheme, asymptotic analysis, orthogonal polynomials, hypergeometric polynomials, special functions.

MSC-class: 33C45 (Primary) 34E05, 33C10 (Secondary)

*e-mail: dominicd@newpaltz.edu

1 Introduction

The Hermite polynomials $H_n(x)$ are defined by [18]

$$H_n(x) = n! \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k}{k!(n-2k)!} (2x)^{n-2k} \quad (1)$$

for $n = 0, 1, \dots$. They satisfy the orthogonality condition [14]

$$\int_{-\infty}^{\infty} e^{-x^2} H_m(x) H_n(x) dx = \sqrt{\pi} 2^n n! \delta_{nm}$$

and the reflection formula

$$H_n(-x) = (-1)^n H_n(x). \quad (2)$$

The Hermite polynomials are special cases of the parabolic cylinder function $U(a, z)$,


$$H_n(x) = 2^{\frac{n}{2}} \exp\left(\frac{x^2}{2}\right) U\left(-n - \frac{1}{2}, \sqrt{2}x\right),$$

which was analyzed by Nico Temme in [5], [19] and [22].

The Hermite polynomials have been extensively studied since the pioneer article of C. Hermite [6] in 1864 (they were previously considered by Fourier and Chebyshev). They have many applications in the physical sciences and are particularly important in the quantum mechanical treatment of the harmonic oscillator [13] (see also [1], [7] and [24] for some extensions). We refer the interested reader to [3] and [15] for further properties and references.

There are several families of orthogonal polynomials which have asymptotic approximations in terms of $H_n(x)$. Some cases studied by Nico Temme include the Gegenbauer [17], Laguerre [12], [16], Tricomi-Carlitz and Jacobi polynomials [10]. He also considered the asymptotic representations of other families of polynomials such as the generalized Bernoulli, Euler, Bessel and Buchholz polynomials in [11].

A rich source of asymptotic relations between the $H_n(x)$ and other polynomials [4], [20], [21] is provided by the Askey-scheme of hypergeometric orthogonal polynomials [8]:



where the arrows indicate limit relations between the polynomials.

In particular, the limit relation between the Charlier polynomials $C_n^{(a)}(x)$ and the Hermite polynomials is given by

$$\lim_{a \rightarrow \infty} (-1)^n (2a)^{\frac{n}{2}} C_n^{(a)} \left(a + x\sqrt{2a} \right) = H_n(x). \quad (3)$$

In this article we investigate the asymptotic behavior of $H_n(x)$ as $n \rightarrow \infty$, using (3) and the asymptotic results on the Charlier polynomials derived in [2]. We believe that our method provides a useful approach to asymptotic analysis and could be used for other families of polynomials of the Askey scheme.

2 Previous results

We define the Charlier polynomials by

$$C_n^{(a)}(x) = {}_2F_0 \left(\begin{matrix} -n, -x \\ - \end{matrix} \middle| -\frac{1}{a} \right), \quad n = 0, 1, \dots \quad (4)$$

with $a > 0$. The following results were derived in [2].

Theorem 1 *As $n \rightarrow \infty$, $C_n^{(a)}(x)$ admits the following asymptotic approximations, with*

$$\Omega^\pm = (\sqrt{n} \pm \sqrt{a})^2. \quad (5)$$

1. $n = O(1)$.

$$C_n^{(a)} \simeq \left(1 - \frac{x}{a}\right)^n \quad (6)$$

2. $x < \Omega^-$, $0 < n < a$.

$$C_n^{(a)} \sim F_3(x) = \exp[\Psi_3(x)] L_3(x), \quad (7)$$

where

$$\Psi_3(x) = x \ln \left(\frac{a+x-n+\Delta}{2a} \right) + n \ln \left(\frac{a-x+n+\Delta}{2a} \right) + \frac{1}{2}(a-x-n+\Delta) \quad (8)$$

and

$$L_3(x) = \sqrt{\frac{a-x-n+\Delta}{2\Delta}}, \quad (9)$$

with

$$\Delta(x) = \sqrt{a^2 - 2a(x+n) + (x-n)^2}. \quad (10)$$

3. $\Omega^+ < x$.

$$C_n^{(a)} \sim F_4(x) = (-1)^n \exp[\Psi_4(x)] L_4(x), \quad (11)$$

where

$$\Psi_4(x) = x \ln \left(\frac{a+x-n-\Delta}{2a} \right) + n \ln \left(\frac{x-a-n+\Delta}{2a} \right) + \frac{1}{2}(a-x-n+\Delta) \quad (12)$$

and

$$L_4(x) = \sqrt{\frac{x-a+n+\Delta}{2\Delta}}. \quad (13)$$

4. $x \approx \Omega^-$, $0 < n < a$.

$$C_n^{(a)} \sim \sqrt{2\pi} \left(\frac{n}{a}\right)^{\frac{1}{6}} (\sqrt{a} - \sqrt{n})^{\frac{1}{3}} \text{Ai} \left[\left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(\Omega^- - x)}{(\sqrt{a} - \sqrt{n})^{\frac{2}{3}}} \right] \times \exp \left[\frac{1}{2}n \ln \left(\frac{n}{a}\right) + x \ln \left(1 - \sqrt{\frac{n}{a}}\right) + \sqrt{an} - \sqrt{n} \right], \quad (14)$$

where $\text{Ai}(\cdot)$ is the Airy function.

5. $\Omega^- < x < \Omega^+$.

$$C_n^{(a)} \sim F_{10}(x) = F_3(x) + F_4(x). \quad (15)$$

6. $x \approx \Omega^+$.

$$C_n^{(a)} \sim \sqrt{2\pi} \left(\frac{n}{a}\right)^{\frac{1}{6}} (\sqrt{a} + \sqrt{n})^{\frac{1}{3}} \text{Ai} \left[\left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(x - \Omega^+)}{(\sqrt{a} + \sqrt{n})^{\frac{2}{3}}} \right] \quad (16)$$

$$\times (-1)^n \exp \left[\frac{1}{2} n \ln \left(\frac{n}{a}\right) + x \ln \left(1 + \sqrt{\frac{n}{a}}\right) - \sqrt{an} - \sqrt{n} \right].$$

3 Limit analysis

From (5) we have, as $a \rightarrow \infty$,

$$\frac{\Omega^\pm - a}{\sqrt{2a}} \rightarrow \pm \sqrt{2n}. \quad (17)$$

Hence, the six regions of Theorem 1 transform into the following regions:

1. Region I: $n = O(1)$.

Setting

$$x = a + \xi \sqrt{2a} \quad (18)$$

in (6) and using (3), we get

$$H_n(\xi) \simeq (2\xi)^n. \quad (19)$$

The formula above is exact for $n = 0, 1$ and is a good approximation when $\xi \gg n$.

2. Region II: $\xi < -\sqrt{2n}$.

From (10) and (18) we have

$$\Delta \sim \sqrt{2}\sigma\sqrt{a} - \frac{\xi n}{\sigma}, \quad a \rightarrow \infty, \quad (20)$$

with

$$\sigma = \sqrt{\xi^2 - 2n}. \quad (21)$$

Using (20) in (8) and (9), we get

$$\frac{n}{2} \ln(2a) + \Psi_3(x) \sim \Phi_1(\xi) \equiv n \ln(\sigma - \xi) + \frac{\xi^2 + \sigma\xi - n}{2}, \quad (22)$$

$$L_3(x) \sim U_1(\xi) = \sqrt{\frac{1}{2} \left(1 - \frac{\xi}{\sigma}\right)}, \quad (23)$$

as $a \rightarrow \infty$. Thus, from (3) we have

$$H_n(\xi) \sim \Lambda_1(\xi) \equiv (-1)^n \exp[\Phi_1(\xi)] U_1(\xi). \quad (24)$$

3. Region III: $\xi > \sqrt{2n}$.

Using (20) in (12) and (13), we obtain

$$\frac{n}{2} \ln(2a) + \Psi_4(x) \sim \Phi_2(\xi) \equiv n \ln(\sigma + \xi) + \frac{\xi^2 - \sigma\xi - n}{2}, \quad (25)$$

$$L_4(x) \sim U_2(\xi) = \sqrt{\frac{1}{2} \left(1 + \frac{\xi}{\sigma}\right)}, \quad (26)$$

as $a \rightarrow \infty$. Hence,

$$H_n(\xi) \sim \Lambda_2(\xi) \equiv \exp[\Phi_2(\xi)] U_2(\xi). \quad (27)$$

Note that

$$\Lambda_1(-\xi) = (-1)^n \Lambda_2(\xi),$$

as one would expect from (2).

4. Region IV: $\xi \approx -\sqrt{2n}$.

Using (18) in (14) we have, as $a \rightarrow \infty$,

$$\frac{n}{2} \ln(2a) + \frac{1}{2} n \ln\left(\frac{n}{a}\right) + x \ln\left(1 - \sqrt{\frac{n}{a}}\right) + \sqrt{an} - \sqrt{n} \sim \Phi_3(\xi),$$

where

$$\Phi_3(\xi) = \frac{n}{2} \ln(2n) - \frac{3}{2}n - \xi\sqrt{2n}. \quad (28)$$

Also,

$$\sqrt{2\pi} \left(\frac{n}{a}\right)^{\frac{1}{6}} (\sqrt{a} - \sqrt{n})^{\frac{1}{3}} \sim n^{\frac{1}{6}}$$

and

$$\left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(\Omega^- - x)}{(\sqrt{a} - \sqrt{n})^{\frac{2}{3}}} \sim -n^{\frac{1}{6}} \sqrt{2} (\xi + \sqrt{2n}).$$

Therefore,

$$H_n(\xi) \sim \Lambda_3(\xi) \equiv (-1)^n \sqrt{2\pi} n^{\frac{1}{6}} \exp[\Phi_3(\xi)] \text{Ai}\left[-n^{\frac{1}{6}} \sqrt{2}(\xi + \sqrt{2n})\right]. \quad (29)$$

5. Region V: $\xi \approx \sqrt{2n}$.

Using (18) in (16) we have, as $a \rightarrow \infty$,

$$\frac{n}{2} \ln(2a) + \frac{1}{2} n \ln\left(\frac{n}{a}\right) + x \ln\left(1 + \sqrt{\frac{n}{a}}\right) - \sqrt{an} - \sqrt{n} \sim \Phi_4(\xi),$$

with

$$\Phi_4(\xi) = \frac{n}{2} \ln(2n) - \frac{3}{2}n - \xi \sqrt{2n}. \quad (30)$$

Also,

$$\sqrt{2\pi} \left(\frac{n}{a}\right)^{\frac{1}{6}} (\sqrt{a} + \sqrt{n})^{\frac{1}{3}} \sim n^{\frac{1}{6}}$$

and

$$\left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(x - \Omega^+)}{(\sqrt{a} + \sqrt{n})^{\frac{2}{3}}} \sim n^{\frac{1}{6}} \sqrt{2} (\xi - \sqrt{2n}).$$

Therefore,

$$H_n(\xi) \sim \Lambda_4(\xi) \equiv \sqrt{2\pi} n^{\frac{1}{6}} \exp[\Phi_4(\xi)] \text{Ai}\left[n^{\frac{1}{6}} \sqrt{2}(\xi - \sqrt{2n})\right]. \quad (31)$$

Once again, we have

$$\Lambda_3(-\xi) = (-1)^n \Lambda_4(\xi)$$

6. Region VI: $-\sqrt{2n} \ll \xi \ll \sqrt{2n}$.

From (15), we immediately obtain

$$H_n(\xi) \sim \Lambda_5(\xi) \equiv \Lambda_1(\xi) + \Lambda_2(\xi).$$

Since $-1 < \frac{\xi}{\sqrt{2n}} < 1$, we set

$$\xi = \sqrt{2n} \sin(\theta), \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}. \quad (32)$$

From (21) we have

$$\sigma = \sqrt{2n} \cos(\theta) i.$$

Thus,

$$n\pi i + \Phi_1(\xi) = \frac{n}{2} [\ln(2n) - \cos(2\theta)] + n \left[\frac{1}{2} \sin(2\theta) + \theta - \frac{\pi}{2} \right] i,$$

$$\Phi_2(\xi) = \frac{n}{2} [\ln(2n) - \cos(2\theta)] - n \left[\frac{1}{2} \sin(2\theta) + \theta - \frac{\pi}{2} \right] i,$$

and

$$U_1(\xi) = \frac{\exp\left(\frac{\theta}{2}i\right)}{\sqrt{2 \cos(\theta)}}, \quad U_2(\xi) = \frac{\exp\left(-\frac{\theta}{2}i\right)}{\sqrt{2 \cos(\theta)}}.$$

Hence,

$$\Lambda_5 \left[\sqrt{2n} \sin(\theta) \right] = \sqrt{\frac{2}{\cos(\theta)}} \exp[\Phi_5(\theta)] \cos(\Theta), \quad (33)$$

with

$$\Phi_5(\theta) = \frac{n}{2} [\ln(2n) - \cos(2\theta)] \quad (34)$$

and

$$\Theta = n \left[\frac{1}{2} \sin(2\theta) + \theta - \frac{\pi}{2} \right] + \frac{\theta}{2}. \quad (35)$$

Using (32), we can write (34) and (35) in terms of ξ

$$\Phi_5(\theta) = \frac{n}{2} [\ln(2n) - 1] + \frac{\xi^2}{2},$$

$$\Theta = \frac{\xi}{2} \sqrt{2n - \xi^2} + \left(n + \frac{1}{2} \right) \arcsin \left(\frac{\xi}{\sqrt{2n}} \right) - n \frac{\pi}{2}.$$

Also,

$$\sqrt{\frac{2}{\cos(\theta)}} = \sqrt{2} \left(1 - \frac{\xi^2}{2n} \right)^{-\frac{1}{4}}.$$

Therefore,

$$\begin{aligned} \Lambda_5(\xi) &= \sqrt{2} \left(1 - \frac{\xi^2}{2n} \right)^{-\frac{1}{4}} \exp \left\{ \frac{n}{2} [\ln(2n) - 1] + \frac{\xi^2}{2} \right\} \\ &\times \cos \left[\frac{\xi}{2} \sqrt{2n - \xi^2} + \left(n + \frac{1}{2} \right) \arcsin \left(\frac{\xi}{\sqrt{2n}} \right) - n \frac{\pi}{2} \right]. \end{aligned} \quad (36)$$

Considering the leading term of (36) as $n \rightarrow \infty$, we obtain

$$\Lambda_5(\xi) \sim \sqrt{2} \exp \left\{ \frac{n}{2} [\ln(2n) - 1] + \frac{\xi^2}{2} \right\} \cos \left(n \frac{\pi}{2} - \xi \sqrt{2n} \right)$$

in agreement with formula (4.14.9) in [9].

4 Zeros

Let us denote by $\zeta_1^n > \zeta_2^n > \dots > \zeta_n^n$ the zeros of $H_n(\xi)$, enumerated in decreasing order. It then follows from (33) that $\zeta_j^n = \sqrt{2n} \sin(\tau_j^n)$, where τ_j^n satisfies

$$n \left[\frac{1}{2} \sin(2\tau_j^n) + \tau_j^n - \frac{\pi}{2} \right] + \frac{\tau_j^n}{2} = \frac{\pi}{2} - j\pi, \quad 1 \leq j \leq n.$$

We can rewrite the equation

$$n \left[\frac{1}{2} \sin(2t) + t - \frac{\pi}{2} \right] + \frac{t}{2} = A$$

as Kepler's equation

$$E - \varepsilon \sin(E) = M, \quad (37)$$

with

$$E = 2t, \quad M = 2 \frac{2A + n\pi}{2n + 1}, \quad \varepsilon = -\frac{2n}{2n + 1}. \quad (38)$$

It is well known [23] that the solution of (37) can be expressed as a Kapteyn series

$$E = M + 2 \sum_{k=1}^{\infty} \frac{1}{k} J_k(k\varepsilon) \sin(kM), \quad (39)$$

where $J_k(\cdot)$ is a Bessel function of the first kind.

Thus, using (38) in (39) with $A = \frac{\pi}{2} - j\pi$, we obtain

$$\tau_j^n = \pi \frac{1 + n - 2j}{2n + 1} + \sum_{k=1}^{\infty} \frac{1}{k} J_k \left(-\frac{2n}{2n + 1} k \right) \sin \left(2\pi \frac{1 + n - 2j}{2n + 1} k \right), \quad (40)$$

for $1 \leq j \leq n$. Using the reflection formula [14] $J_k(-x) = (-1)^k J_k(x)$, we can write (40) as

$$\tau_j^n = \frac{\pi}{2} - \frac{\pi}{2} (4j-1) N^{-1} - \sum_{k=1}^{\infty} \frac{1}{k} J_k \left[(1 - N^{-1}) k \right] \sin \left(\frac{4j-1}{N} k \pi \right), \quad (41)$$

where $N = 2n + 1$.

References

- [1] M. Aunola. The discretized harmonic oscillator: Mathieu functions and a new class of generalized Hermite polynomials. *J. Math. Phys.*, 44(5):1913–1936, 2003.
- [2] D. Dominici. Asymptotic analysis of the Askey-scheme I: from Krawtchouk to Charlier. Preprint, arXiv: math.CA/0501072, 2005.
- [3] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vols. I, II*. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1953.
- [4] C. Ferreira, J. L. Lopez, and E. Mainar. Asymptotic relations in the Askey scheme for hypergeometric orthogonal polynomials. *Adv. in Appl. Math.*, 31(1):61–85, 2003.
- [5] A. Gil, J. Segura, and N. M. Temme. Integral representations for computing real parabolic cylinder functions. *Numer. Math.*, 98(1):105–134, 2004.
- [6] C. Hermite. Sur un nouveau développement en série de fonctions. *Compt. Rend. Acad. Sci. Paris*, 58:93–100, 1864.
- [7] R. Hinterding and J. Wess. q -deformed Hermite polynomials in q -quantum mechanics. *Eur. Phys. J. C Part. Fields*, 6(1):183–186, 1999.
- [8] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue. Technical Report 98-17, Delft University of Technology, 1998. <http://aw.twi.tudelft.nl/~koekoek/askey/>.

- [9] N. N. Lebedev. *Special functions and their applications*. Dover Publications Inc., New York, 1972.
- [10] J. L. López and N. M. Temme. Approximation of orthogonal polynomials in terms of Hermite polynomials. *Methods Appl. Anal.*, 6(2):131–146, 1999. Dedicated to Richard A. Askey on the occasion of his 65th birthday, Part II.
- [11] J. L. López and N. M. Temme. Hermite polynomials in asymptotic representations of generalized Bernoulli, Euler, Bessel, and Buchholz polynomials. *J. Math. Anal. Appl.*, 239(2):457–477, 1999.
- [12] J. L. López and N. M. Temme. Convergent asymptotic expansions of Charlier, Laguerre and Jacobi polynomials. *Proc. Roy. Soc. Edinburgh Sect. A*, 134(3):537–555, 2004.
- [13] L. Pauling and E. B. Wilson, Jr. *Introduction to quantum mechanics*. Dover Publications Inc., New York, 1985.
- [14] J. Spanier and K. B. Oldham. *An Atlas of Functions*. Hemisphere Pub. Corp., 1987.
- [15] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition, 1975.
- [16] N. M. Temme. Asymptotic estimates for Laguerre polynomials. *Z. Angew. Math. Phys.*, 41(1):114–126, 1990.
- [17] N. M. Temme. Polynomial asymptotic estimates of Gegenbauer, Laguerre, and Jacobi polynomials. In *Asymptotic and computational analysis (Winnipeg, MB, 1989)*, volume 124 of *Lecture Notes in Pure and Appl. Math.*, pages 455–476. Dekker, New York, 1990.
- [18] N. M. Temme. *Special functions*. A Wiley-Interscience Publication. John Wiley & Sons Inc., New York, 1996.
- [19] N. M. Temme. Numerical and asymptotic aspects of parabolic cylinder functions. *J. Comput. Appl. Math.*, 121(1-2):221–246, 2000. Numerical analysis in the 20th century, Vol. I, Approximation theory.

- [20] N. M. Temme and J. L. López. The role of Hermite polynomials in asymptotic analysis. In *Special functions (Hong Kong, 1999)*, pages 339–350. World Sci. Publishing, River Edge, NJ, 2000.
- [21] N. M. Temme and J. L. López. The Askey scheme for hypergeometric orthogonal polynomials viewed from asymptotic analysis. *J. Comput. Appl. Math.*, 133(1-2):623–633, 2001. Proceedings of the Fifth International Symposium on Orthogonal Polynomials, Special Functions and their Applications (Patras, 1999).
- [22] N. M. Temme and R. Vidunas. Parabolic cylinder functions: examples of error bounds for asymptotic expansions. *Anal. Appl. (Singap.)*, 1(3):265–288, 2003.
- [23] G. N. Watson. *A treatise on the theory of Bessel functions*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995.
- [24] A. Zarzo, J. S. Dehesa, and J. Torres. On a new set of polynomials representing the wave functions of the quantum relativistic harmonic oscillator. *Ann. Numer. Math.*, 2(1-4):439–455, 1995. Special functions (Torino, 1993).